#### **Open Access**

#### **RESEARCH PAPER**

# ASSESSMENT OF POST STROKE DISABILITY AND ITS MANAGEMENT USING STRENGTHENING EXERCISE

## Md. Ruhul Amin<sup>\*1</sup>, Md. Redwanur Rahman<sup>1</sup>, Md. Moniruzzaman Sarker<sup>2</sup>and Md. Nurul Islam<sup>3</sup>

<sup>1</sup>Institute of Environmental Science, University of Rajshahi, Bangladesh

<sup>2</sup> Department of Zoology, University of Rajshahi, Bangladesh

<sup>3</sup> Department of Medicine, Rajshahi Medical College and Hospital, Bangladesh

**Keywords:** Stroke, Glasgow Coma Scale, Barthel Disability Index, Asian Stroke Disability Scale and Strengthening Exercise

Publication date: February 14, 2023

#### Abstract

Stroke is a focal episode of neurological deficit, brain tissue damage and the leading cause of death and disability. A cross-sectional analytical study was conducted to identify the disability on stroke patients, second fold of the study was experimental, and it was conducted to evaluate the efficacy of strengthening exercise on physically disabled of stroke patients. A pretested, semi-structured standard questionnaire and randomized sampling technique used to collect the data, the number of calculated sample size was 105, and SPSS version 20.0 was used for data analysis. Study periods were the 2 years duration from July 2018 to 30<sup>th</sup> June 2020. Study found that average age of the respondents was $55.68 \pm 12.77$  years. Among the respondents, 58.10% were male and 41.90% were female. Study revealed that mean Barthel Disability Index had 29.71  $\pm$  27.49 and mean Asian Stroke Disability Scale had 5.76  $\pm$ 2.15. However, patients were suffering from physical, speaking, psychological, visual and auditory disability. There were statistically significant findings after intervention of strengthening exercise where t-value were -17.737 and 12.464, which measured by Barthel Disability Index and Asian Stroke Disability Scale. A comprehensive rehabilitation program includes muscle-strengthening exercise techniques were statistically significant effects on physical disability of stroke patients.

\*Corresponding Author: Md. Ruhul Amin\*, Email: ruhul31physio@ yahoo.com

#### Introduction

Stroke is a disease of the brain where there is sudden onset of mostly focal lesion due to occlusion or rupture of a cerebral blood vessel and its symptomatology should last for more than twenty four hours. The incidence of stroke is increasing in this country in comparison to developed country. In our country the prevalence of stroke above the age of 40 is 370/100000 (Mohammad et al., 2011).Stroke is the leading cause of death and disability in Bangladesh, only a few government and non-governmental organizations provide rehabilitation for stroke patients into the Bangladesh health system (Mamin et al., 2017).Oral contraceptive pill, pregnancy, connective tissue diseases with vacuities, hematological variables. drug abuse. smoking, congenital heart diseases, family history and some genetic diseases were responsible for stroke (Kristensen et al., 1997).In Asia, the problem of stroke has a particularly strong impact, more than half of the world's population lives in Asia, but stroke is the predominant vascular disease in many parts of Asia (Shi et al., 1998). The prevention, identification and control of the cardiovascular risk factors at individual and population level is poor in Asia because of ignorance and high cost of the interventions so diseases burden is likely to increase in the near future(Taylor et al., 1996). In a hospital based study, it was revealed that hypertension is the most important risk factor for stroke in Bangladesh (Rahman, 2002).History of hypertension, diabetes mellitus, central obesity, Psychosocial stress, smoking were the risk factors of ischemic stroke, whereas alcohol intake were significant risk factors for hemorrhagic stroke (Donnel et al., 2010). High blood pressure,

smoking, severe psychological stress. atrial fibrillation, previous transient ischemic attacks. and intermittent claudication were the independent risk factors for hemorrhagic stroke non (Harmsen et al., 1990). First-degree relative and genetic risk factors may be important in the incidence of intracerebral hemorrhage (Woo et al., 2002). Increased risk of intracerebral hemorrhage with prior cerebral infarction, and these data suggest that this occurs independent of aspirin or anticoagulant use (Gorter, 1999).

Stroke is also in the top five diseases with the greatest burden of disease, based on disability (Loo and Gan, 2012).Stroke causes long-term disability resulting from secondary complications common after a stroke such as heart disease, bedsore, urinary tract infection, respiratory infection (Eng and Tang, 2007). Physiotherapy plays a significant role in the rehabilitation of patients' functional movements after stroke. Early mobilization, balance, muscle strengthening exercise, endurance exercise, gait training, hydrotherapy and social activation of patients or his family (Macek et al., 2020). Cardiorespiratory training, therapeutic exercise, task-oriented training, mirror therapy, neuromuscular electrical stimulation shows promise as an intervention for stroke(Lin and Dionne, 2018). Or ofacial therapy as an emphasis of movement of tongue and hyoid muscles. (Konecny et al., 2017).Constraint-induced movement therapy, mental practice. intervention for sensory impairment, these may be effective intervention (Pollock et al., 2014). However, functional approach, Bobath approach and motor learning approach had statistically significant improvements their disability (HoChung,

2014). Physiotherapy usually provided in the first months after stroke is effective but in the chronic phase is uncertain (Ferrarello and Toro, 2011).Orthopedic principle, motor learning or mixed methods of these treatment principles (Pollock et al., 2008). Whereas functional training of the upper limb movement therapy, treadmill training with or without body weight support (Van Peppen et al., 2004). More than half of the patients who survive the first month after a stroke will require specialized rehabilitation (Pollack and Disler, 2002).The study was conducted to identify the disability of stroke patients and assess the efficacy of strengthening exercises on physically disabled stroke patients.

#### 2. Materials and Methods

#### 2.1 Study Populations

The study populations were the patients who were suffering different types of stroke and admitted in the National Institute of Neuro Sciences & Hospital.

#### 4.2 Study Area and Study Sites

This study was conducted in Dhaka city. This study was employed among the patients admitted in National Institute of Neuro Sciences & Hospital (NINSH) in Dhaka city. This study sites were in NINSH. Data were collected from selected words (one male and one female) from NINSH.

#### 2.3 Study Design and Study Periods

The present study was cross-sectional analytical study and this study was the most feasible for identifying the disability on stroke patients, and second fold of the study was experimental study that helps to explore the efficacy of strengthening exercise on physically disabled of stroke patients. These study periods were the two years duration from 1<sup>st</sup> July 2018 to 30<sup>th</sup> June 2020.

## **2.4 Data Collection Tools, Instruments, and Sampling Techniques**

A pretested, semi-structured questionnaire and randomized sampling technique was applied to collect the data with a sample size was 105. One male and one female wards were select for collecting sample, 61 males and 44 females patients were included the study among them 30 patients were included for experimental group and another 30 patients were included for control group.

#### 2.5 Inclusion and Exclusion Criteria

Those who are willing to give consent and participate for interview, irrespective of sex and clinically diagnosed stroke patients. Patients or attendants disagree to Participate in the study.

#### 2.6 Calculation of Sample Size

Following formula is be used to determine the sample size.

$$n = \frac{z^2 p q}{d^2}$$

Here

n = the desired sample size

z = the standard normal deviate usually set at 1.96 which correspondents to 95% confidence level

p = 6.59%(Estimated prevalence of Stroke in Philippines) (Looand Gan, 2013)

$$q = 1 - p = 1.00 - 0.0659$$

q = 0.9341

d = degree of accuracy desired, usually set at 0.05%.

Now, required sample size

$$n = \frac{(1.96)^2 \times 0.0659 \times 0.9341}{(0.05)^2}$$
$$= 94.59$$
$$= 95$$

When 90% response rate, so sample size = 95+10% of 95

- = 95+9.5
- = 104.5
- = 105

So required sample size was 105.

#### 2.7 Data Collection Technique

From the participant by face-to-face interview or attendants, socio-economic information's were collected by interview methods. Stroke related and Physiotherapy treatment and outcome related information detected by history. were physical examination and medical records if possible. Respondents' mobile numbers were included on questionnaire. Data were collected from survey of the patients those who clinically diagnosis stroke patients through questionnaire by personal interview.

#### 2.8 Therapeutic Management

Effective rehabilitation on a coordinated activity of multidisciplinary teamwork of early mobilization and therapeutic exercise especially muscle strengthening techniques were applied for intervention.

#### 2.9 Measurement Scales

Measurement of disability of the respondents of both group (experimental and control) used by Barthel index and the Asian stroke disability scale.

### 2.10 Data Processing and Analysis

After collection of data, it was checked and rechecked thoroughly with competently and these data were analyzed by using SPSS software version 20.After analyzed the data set all information were interpreted and presented tabulated form (univariate and bivariate), grapes (bar diagram and pie chart) and pared t-test was applied for both experimental and control groups.

#### 2.11 Informed Consent

There was no human risk, no medication and no invasive procedure only exercise programs were done for intervention. Informed consent was taken before starting the interview of the respondents, all information was kept confidential, used only research work. Respondents had right to refuse and withdraw from the research work.

## 2.12 Ethical Clearance

The research proposal approved by the ethical committee of National Institute of Neuro-Sciences and Hospital, Shere-E-Bangla Nagar, Dhaka, Bangladesh.

#### 3. Results

 Table 01: Socio-demographic Information (n=105)
 Information (n=105)

Variables	Items	Frequency	Percentage
	≤ 30	2	1.90
	31-40	13	12.40
	41-50	25	23.80

	51-60	26	24.80
Age in Years	61-70	33	31.40
	≥71	6	5.70
	Total	105	100.00
	Mean ± SD		55.68 ± 12.77
	Male	61	58.10
Gender	Female	44	41.90
	Total	105	100.00
	Lower class	04	3.80
	Lower middle class	57	54.30
Social Classes	Upper middle class	38	36.20
	Upper class	06	5.70
	Total	105	100.00
	Mean ± SD	277	57.14 ± 21080.67

Table 01 shows that more than 85.00% of the respondents belonged to their age group were  $\geq 40$  years with their mean age was 55.68  $\pm$  12.77 years. Study reveals that 58.10% of the respondents were male and 41.90% of the respondents were female. Study found that 3.80%, 54.30%, 36.20% and 5.70% of the respondents were lower class, lower middle class, upper middle class and upper class respectively with their mean monthly income was 27757.14  $\pm$  21080.67 BDT.



Figure 01: Distribution of the study subjects by Hypertension.

Figure 01 reveals that 35.20% of the respondents suffered systolic hypertension, 30.50% of the respondents suffered diastolic hypertension and 34.30% of the respondents had normal pressure.

Variables	Items	Frequency	Percentage
Barthel Index	$\leq 60$ (Institutionalized)	89	84.80
	61-100 (Able to live independently)	16	15.20
	$\geq$ 101 ( Capable of carried himself)	00	00
Total		105	100.0
	Mean ± SD	$29.71 \pm 27.49$	
The Asian Stroke	$\leq$ 3 (Live independently)	15	14.30
Disability Scale	≥4 (Need Institutionalized)	90	85.70
	Total		100.0
	Mean ± SD	5.76±	2.15

Table 02: Distribution of the study subjects by disability measurement of post stroke patients (n=105).

Table 02 shows that 84.80% of the respondents need institutionalized support for rehabilitation and 15.20% of the respondents able to live independently with their mean disability was  $29.71 \pm$ 27.49. It was measured by Barthel Index. Study showed that 14.30% of the respondents live independently and 85.70% of the respondents need institutionalized for rehabilitation with their mean disability was  $5.76 \pm 2.15$ . It was measured by the Asian Stroke Disability Scale.



Figure 02: Distribution of the study subjects by types of stroke (n=105).

Figure 02 shows that 29.50% of the respondents suffered ischemic stroke, 64.80% of the respondents suffered hemorrhagic stroke and 5.70% of the respondents suffered recurrent stoke.

Glasgow Coma Scale	Frequency	Percentage	
$\leq 8$ (Severe injury)	41	39.00	
9-12 (Moderate injury)	40	38.10	
13-14 (Mild injury)	11	10.50	
15 (Normal or healthy)	13	12.40	
Total	105 100.00		
Mean ± SD	$10.05 \pm 3.00$		

Table 03 reveals that 39.00%, 38.10%, 10.50% and 12.40% of the respondents' severity of brain injury had severe, moderate, mild and normal respectively with their mean GCS was  $10.05 \pm 3.00$  and it was measured by Glasgow Coma Scale.



Figure 03: Distribution of the study subject by types of disability.

Figure 03 shows that 74.30%, 65.70%, 4.80%, 6.70% and 4.80% of the respondents had physical, speaking, psychological, visual, and auditory disability respectively.

Table 04: Distribution of the study subjects by findings of computerized tomography scan (n=105).

CT Scan Findings	Frequency	Percentage
Ischemic changes	25	23.80
Intracerebral hemorrhage	56	53.30

Sub arachnoid hemorrhage	19	18.10
Recurrent- First infraction then hemorrhage	03	2.90
Recurrent- First ischemic then ischemic	01	1.00
Recurrent- First hemorrhage then hemorrhage	01	1.00
Total	105	100.00

Table 04 shows that 23.80%, 53.30%, 18.10%, 2.90%, 1.00% and 1.00% of the respondents Ct scan Findings were ischemic changes, in tracerebral hemorrhage, sub arachnoid hemorrhage,

(recurrent) first infraction then hemorrhage, (recurrent) first ischemic then ischemic and(recurrent) first hemorrhage then hemorrhage respectively.

Table 05: Distribution of respondents by efficacy of strengthening exercise technique (n=40+40).

Variables	Group	Mean		t-Value	p-Value
Barthel Index	Study Group	Pre test	Post test		
	Study Gloup	$30.87 \pm 21.33$	$65.25 \pm 19.99$	-17.737	0.000
	Control Group	Pre test	Post test	1 820	
	Control Oloup	$26.87 \pm 25.66$	$28.95 \pm 25.06$	-1.829	0.075
The Asian Stroke	Study Group	Pre test	Post test		
Disability Scale	Study Oloup	$5.45 \pm 1.60$	$3.17 \pm 1.27$	12.464	0.000
2	Control Group	Pre test	Post test	1 863	
	Control Group	$6.12 \pm 1.93$	$5.85 \pm 1.90$	1.005	0.070

#### Paired t-test

Table 05 shows that, t-value was -17.737 which were greater than tabulated t-value 1.697 and (p=0.000>0.05) that was statistically significant findings of study group which was measured by Barthel Index of Activity. Compare to control group, t-value was -1.829 which were greater than tabulated t-value 1.697 and

(p=0.075>0.05) that was statistically not significant findings which was measured by Barthel Index of Activity. Study revealed that t-value was 12.464and (p=0.000>0.05) which were greater than tabulated t-value 1.697 that was statistically significant findings of study group which was measured by the Asian Stroke Disability Scale.

Table 06: Distribution of the study subjects by association between types of stroke and stroke disability of the respondents.

Types of	Stroke Disability		Total	$\chi^2$	p-value
stroke	≤3	$\geq 4$			
Ischemic	1(1)	30(28.6)	31(29.5)	6.303	0.021
Hemorrhagic	14(13.3)	54(51.4)	68(64.8)		
Recurrent	0(0)	6(5.7)	6(5.7)		
Total	15(14.3)	90(85.7)	105(100)		

Results were expressed as number (%),  $\chi^2$  experiment was conducted and p<0.05 was level of significance

significant association between types of stroke and disability of the respondents of the respondents (p=0.021<0.05) and it was measure by Asian Stroke Disability Scale.

Table 06 shows that there was a statistically

Variables	Variables	r-value	p-value
The Asian Stroke	Body mass index	-0.321	0.001
Disability Index	Glasgow coma scale	-0.318	0.001
	Birthel Index	-0.818	0.000
Birthel Index	Body mass index	0.218	0.025
Dif thei maex	Random blood sugar	0.236	0.026
	Glasgow coma scale	0.392	0.000

Table 07: Distribution of the respondents by Pearson correlation of level of disability with tabulated variables.

p- value obtained from Pearson correlation test

Table 07 shows that there were negative correlations of the Asian Stroke Disability Index with Body mass index (p= 0.001>0.05 and r= -0.321), Glasgow coma scale (p= 0.001 > 0.05 and r= -0.318) and Birthel Index (p= 0.000>0.05 and r= -0.818). While positive correlation of Birthel Index with Body mass index (p= 0.025>0.05 and r=0.218), Random blood sugar (p= 0.026>0.05 and r=0.236) and Glasgow coma scale (p= 0.000>0.05 and r=0.392). Study revealed that the Asian Stroke Disability Index negatively influences on Birthel Index, Body mass index and Glasgow coma scale. Whereas, Birthel Index positively influences on Body mass index, Random blood sugar and Glasgow coma scale. These findings were statistically significant.

#### 4. Discussions

Study found that one third of the respondents age were 61 to 70 years, second age group were 41-50 years and it was the 24.80% of the respondents, another age group were 31-40 years and it was 23.80%, rest of them were 19.10% and these age group were above 71 years and below 30 years. These findings were nearly similar findings to the study carried out by Wu *et al.* in 2010. Mean age of the

study subjects was 55.68  $\pm$  12.77 years, where mean age of the respondents of review were  $53 \pm 16$  years and these findings were closely similar findings to the study carried out by Komolafe et al. in 2015. Among the respondents, 58.10% were male and 41.90% were female, where another similar type study found that 46.2% of the patients were male and 53.8% of the patients were female(Komolafe et al., 2015).Study revealed that 3.80%, 54.30%, 36.20% and 5.70% of the respondents were lower class, lower middle class, upper middle class and upper class respectively with their mean monthly income was 27757.14  $\pm$  21080.67 BDT. Opposite findings of the study carried out by Wang et al. in 2019 and revealed that upper class people were more suffering of stroke (Wang et al., 2019). Of the respondents, 20.00% lived in urban and 80.00% lived in rural area. Another similar type study conducted by Wu et al. in 2019 and found that rural people of China were more affected that urban people (Wu et al., 2019).

Study revealed that 29.50% of the respondents suffered ischemic stroke, 64.80% suffered hemorrhagic stroke and 5.70% suffered recurrent stoke. Another dissimilar finding to the study conducted by Momen *et al.* in 2016 and revealed that 66% were ischemic stroke and 34% were

hemorrhagic. Study revealed that 35.20% of the respondents suffered systolic hypertension, 30.50% suffered diastolic hypertension, and 34.30% had normal pressure. Similar type study carried out by Biderafsa et al., in 2015 and showed that systolic blood pressure were 3.5% and diastolic blood pressure were 4.87% in total burden of stroke risk (Biderafsa et al., 2015).Mean Barthel disability index was29.71  $\pm$  27.49 and mean Asian Stroke Disability Score was 5.76  $\pm$  2.15. These findings were similar findings to the study carried out by Pinedo et al. in 2017 and Dubuc et al. in 2015. There was a statistically significant finding of study group after intervention of different types of muscle strengthening exercise which was measured by Barthel Index of Activity. These findings were similar findings to the study carried out by Macek et al. 2020 and revealed that muscle is strengthening exercise. endurance exercise, cyclic bilateral arm and leg training, gait training had significant effect on stroke disability (Macek et al., 2020). There was another statistically highly significant finding of study group after intervention of different types of muscle strengthening exercise which was measured by the Asian Stroke Disability Scale. Another similar type study carried out by Konecny et al. in 2017.

#### 5. Conclusion

Study concludes that assessment of disability by using of Barthel Index of Activity and Asian Stroke Disability Scale and found that most of the stroke patients need institutional support for better management. However, this physical disability is managed by using muscle strengthening exercise technique and it was statistically significant. Effective rehabilitation on a coordinated activity of different types of therapeutic exercises were the components of intervention.

#### 6. Recommendations

Based on the review findings, the following recommendations made to prevent and reduce risk of stroke.

- 1. Raising awareness among the population about the disability of stroke patients and its severity.
- 2. Continuous health education program by using print media and electronic media about the rehabilitation of stroke patients.
- 3. Further study is needed to assess the effective management of modern physiotherapy intervention for minimizing physical, social and psychological disability.

#### References

Biderafsa A, Karami M, Faradmal J and Poorolajal J. 2015. Estimated the potential impact fraction of hypertension as the main risk factor of stroke: application of the distribution shift method. *Journal of Epidemiology* and Global *Health*,**5(3)**:231-33.

Boysen G,Nyboe J, Appleyard M, Sorensen PS, Boas J,Somnier F,Jensen G and Schnohr P.1988. Stroke incidence and risk factors for stroke in Copenhagen, Denmark.*Strok*,.**19**(**11**):1345–53.

Donnel MJ, Xavier D, Liu L, Zang H, Chin SL, Melacini PR. 2010. Risk factors for ischemic and intracerebral hemorrhagic stroke in 22 countries (the inter stroke study): a case-control study. *The Lancet*,**376**(**9735**):112-23.

Dubuc V, Choi P, Hill MD and Coutts SB. 2015. Can 90-day NIHS be used for

outcome assessment in TIA and minor stroke studies? *Cerebrovasc Dis*,40(1-2):97-98.

Eng JJ and Tang PF. 2007. Gait training strategies to optimize walking ability in people with stroke: a synthesis of the evidence. *Expert Rev Neurother*, **7(10)**:1417-36.

Ferrarello F and Toro J. 2011. Efficacy of physiotherapy interventions late after stroke: a meta-analysis. *J Neurol Neurosurg Psychiatry*,**82**:136-43.

Gorter JW. 1999. Major bleeding during anticoagulation after cerebral ischemia: patterns and risk factors: Stroke Prevention in Reversible Ischemia Trial (SPIRIT): European Atrial Fibrillation Trial (EAFT) Study Groups. *Neurology*, **53**:1319–127.

Harmsen P, Rosengren A, Tsipogianni A and Wilhelmsen L. 1990. Risk factors for stroke in middle-aged men in Goteborg, Sweden. *Stroke*, **21**(**2**):223–29.

HoChung BP. 2014. Effect of different combinations of physiotherapy treatment approaches on functional outcomes in stroke patients: a retrospective analysis. *Hong Kong Physiotherapy Journal*, **32(1)**:21-27.

Komolafe MA, Olaogu M, Adebiyi AM, Obembe AO, Fawale MB and Adebowale AA. 2015. Stroke risk factors among participants of a world stroke day awareness program in South-Western Nigeria. *Nigerian Journal of Clinical Practic*, **18(6)**:807-09.

Konecny P, Elfmark M, Bastolova P and Gaul-Alacova P. 2017. New orofacial Physiotherapy of dysphagia after stroke. *Int J Phys Med Rehabil*,**5**(**3**):406-10.

Kristensen B, Malm J, Carberg B, Stegmayr B, Backman G, Fagerlung M. 1997. Epidemiology & etiology of ischemic Stroke in young adults aged 18-44 Yrs in Northern Sweden. *Stroke Bangladesh*,**28**:1702-09.

Lin SH and Dionne T. 2018. Intervention to improve movement and functional outcomes in adult stroke rehabilitation: review and evidence summery. *J Participat Med*,10910:03-16.

Loo KW and Gan SH. 2012. Burden of stroke in Malaysia. *Int J Stroke*, **7**(2):165-67.

Loo KW and Gan SH. 2013. Burden of stroke in the Philippines. *Int J Stroke*, **8(2)**:131-4.

Macek Z, Kolar M, Tucic M and Mandic M. 2020. Recommendations for Physiotherapy Intervention After Stroke. *Ann Physiother Clin*, **2(1)**:1011-20.

Mamin FA, Islam MS., Rumana FS andFaruqui F.2017. Profile of stroke patients treated at a rehabilitation centre in Bangladesh. *BMC Research Notes*, **10**:520.

Mohammad QD, Habib M, Hoque A. 2011. Prevalence of stroke above forty years. *Mymensingh Med J*, **20** (**4**):640-44.

Momen TF, Lakhair MA, Shaikh M, Raque A and Rind MS. 2016. Sociodemographic risk factors for hemorrhagic and ischemic stroke: a study in tertiary care hospital of Hyderabad. *Pakistan Journal of Neurological Science(PJNS,* **11(1)**:24-29.

Pinedo S, SanMartin V, Zaldibar B, Miranda M, Tejada P, Erazo P. 2017. Quality of life of stroke patients and their caregivers. *Int J Phys Med Rehabil*, **5**(1):386-72.

Pollack MRP and Disler PB. 2002. Rehabilitation of patients after stroke. *Med J*,**177(8)**:452-56. Pollack MRP and Disler PB. 2002. Rehabilitation of patients after stroke. *Med J*,**177(8**):452-56.

Pollock A, Baer GD, Langhorne P and Pomeroy VM. 2008. Physiotherapy treatment approaches for stroke. *Stroke*,**39(2)**:519-20.

Pollock A, Farmer SE, Brady MC, Langhorne P, Mead GE, Mehrholz J and Vanwijck F. 2014. Intervention for improving upper limb function after stroke. *Cochrane Database Syst Rev*,**11**: 01-140.

Rahman KM. 2002. Risk factors & clinical presentations--a study of eighty-five hospital admitted stroke cases. Mymensingh Med J,11(2):113-15.

Shi F, Hart RG, Sherman DG and Tegeler CH. 1998. Stroke in the People's Republic of China. *Strok*,**20**:1581–85.

Talhatu HK and Funmilayo FO. 2007. Balance retraining in post stroke patients using a simple, effective and affordable technique. *AJNS*, **26**(**2**):39-47.

Taylor TN, Davis PH, Torner JC, Holmes J, Meyer JW and Jacobson MF. 1996. Lifetime cost of stroke in the United States. *Stroke*, **27**:1459-66.

Van Peppen RP, Kwakkel G, Wood-Dauphinee S, Hendriks HJ, Van der Wees PJ and Deeker J. 2004. The impact of Physical therapy on functional outcomes after stroke: what's the evidence? *Clinical Rehabilitation*,**18**(**18**):833-62.

Wang S, Shen B, Wu M et al. 2019. Effects of socioeconomic status on risk of ischemic stroke: a case-control study in Guangzhou population. *BMC*. 19:648.

Wo D, Laura R, Sauerbeck RN, Brett M, Kissela MD, Jane C et al. 2002. Genetic

and Environmental Risk Factors for Intracerebral Hemorrhage Preliminary Results of a Population-Based Study. *Stroke*,**33**:1190-96.

Wu CY, Wu HM, Lee JD and Weng HH. 2010. Stroke risk factors and subtypes in different age groups: a hospital- based study. *Neurol India*,**58**(**6**):863-68.

Wu Y, Fan Z, Chen Y, Ni J, Liu J, Han J, Ren L, Tu J, Ning X and Wang J. 2019. Determinants of developing stroke among low-income, rural residents: a 27-years population-based, prospective cohort study in Northern China. *Front. Neurol*, **10**:57-65.