



Evaluation of Antimicrobial Activity of Some Medicinal Plants in Bangladesh

Mrityunjoy Das^{1*}, Mahfuzur Rahman², Shyam Sundar Shaha³ and Md. Belal Uddin^{*}

¹ Department of Microbiology, GonoBishwabidyalay, Savar, Dhaka, Bangladesh

² Geography and Environmental Studies, University of Rajshahi, Bangladesh

³ Department of Biochemistry and Molecular Biology, GonoBishwabidyalay, Savar, Dhaka, Bangladesh.

* Department of Biochemistry & Molecular Biology, University of Rajshahi, Bangladesh.

ARTICLE INFO

Keywords:

Ocimumtenniflorum,
Centellaasiatica,
Euphorbia hirta,
antimicrobial activity,
inhibition zone.

ABSTRACT

Ethanollic extract of three medicinal plants namely Ocimumtenniflorum, Centellaasiatica and Euphorbia hirta were evaluated for their therapeutic potential as antimicrobial agent against six standard organisms three bacteria and threefungi, using paper disc method. The antimicrobial effectiveness of callus cultures of E. hirta with inhibition zone of 18 mm was found highest against C. gramineum and in O. tenniflorum with inhibition zone of 28 mm was found highest against K. pneumonia. On the Contrary, in in vivo analysis of antimicrobial efficacy of O.tenniflorum with an inhibition zone of 16 mm was found to the highest against P. chrysogenum and E. hirta with an inhibition zone of 24 mm was found to the highest against K. pneumonia.

1. Introduction

Medicinal plants have been used since ancient times to treat illness. Plants used as raw materials in herbal medicine are called medicinal plants (Mohammed 2019). The U.S. Forest Service estimates that 40% of pharmaceutical drugs used in the Western world are derived from plants (Medical Botany 2023). Seven thousand medical compounds are derived from plants in the modern pharmacopeia. Herbal medicine combines traditional empirical knowledge with modern science (Mohammed 2019). A medicinal plant is considered an important source of prevention against various diseases (Mohammed 2019). The essential medicine component is extracted from different parts of the plants. (Kumar *et al.*, 2017). In underdeveloped countries, people use medicinal plants as a substitute for medicines. There are different species of plants around the world. Herbs are one of them, which have different shapes, colors, and leaves (Wu *et al.*, 2007).

Traditional uses of plants have led to investigating their bioactive compounds through screening programs, which have resulted in the detection of a significant number of therapeutic properties.

The search for plants with antimicrobial activity have taken on increasing importance in recent years, due to growing global concern over the alarming increase in the rate of infection with antibiotic-resistant microorganisms or multi-resistant microbes. Many studies have been conducted with extracts from various plants, testing antimicrobial activity as well as for the discovery of new antimicrobial compounds (Dygeraket al., 2002, Ibrahim et al., 2002, Bassam et al., 2006). Plants are used medicinally in different countries and are a source of many potent and powerful drugs (Sasidharan et al., 1998, Semraet al., 2006, Suffredini et al., 2006, Kubmarawa et al., 2007). Interest in scientific investigation of O. tenniflorum, C. asiatica and E. hirta is based on claims of its effective use for the treatment of many diseases.

Corresponding author.

E-mail address: dr.mrityunjoydas@gmail.com (Mrityunjoy Das), drmbuddin@gmail.com (Md. Belal Uddin)

Receive 01 July 2024; Revised Received 27 October 2024; Accepted 11 November 2024;

Available Online 12 December 2024;

Published by Institute of Environmental Science, University of Rajshahi.

The present paper discusses in vitro and in vivo screening of medicinal plants of Bangladesh for their antimicrobial activity. Therefore, this research regarding the antimicrobial activity of these plants is expected to enhance the use of *O. tenniflorum*, *C. asiatica* and *E. hirta*, diseases caused by the test against the pathogens. Pathogenic organisms were selected for study based on their clinical and pharmaceutical importance as well as for their potential to cause contamination of food and drugs.

2. Materials and Methods

2.1. Plant material

Plant materials were collected from Rajshahi university campus Rajshahi, Bangladesh.

2.2. Preparation of plant extract

Fifty gm. powder of each plant material (aerial part of the plant and callus) was percolated with 250 ml of 80% ethanol many soxhlet and the resulting residue was evaporated to dryness to give a residue, which was then used to perform antimicrobial assay.

2.3. Test Microorganism

Pure culture of all the bacteria, namely *K. pneumoniae*, *S. dysenteriae*, *E. coli*, were obtained courtesy of the Department of Microbiology, GonoBishwabidyalay, Savar, Dhaka, Bangladesh, while the fungi tested namely *A. niger*, *C. gramineum*, *P. chrysogenum* were obtained University of Rajshahi, Rajshahi, Bangladesh by the kind permission of the Department of Botany, which were preserved on nutrient broth and potato dextrose agar (PDA) respectively.

2.4. Antimicrobial assay

Antimicrobial activity of the extracts was tested using the filter paper disc method” (Gould *et al.*, 1952).

3. Results and Discussion

Many possible sources of antibiotic extraction and synthesis of antibiotics have been extensively developed, but the search for a better, safer, and more economic source is still necessary. In this context, attempts have been made to sort various crude extract to identify their antimicrobial potential. In the present investigation ethanolic extract of *O. tenniflorum*, *C. asiatica* and *E. hirta* were screened for antimicrobial activity.

Table 1. Anti-microbial activities of crude ethanolic extract of *O. tenniflorum*, *C. asiatica* and *E. hirta*

Plant extract tested	Test method	Zone of inhibition (in mm) against test organisms					
		Fungi			Bacteria		
		<i>A. niger</i>	<i>C. gramineum</i>	<i>P. chrysogenum</i>	<i>K. pneumoniae</i>	<i>S. dysenteriae</i>	<i>E. coli</i>
<i>O. tenniflorum</i>	In vivo	10	10	16	20	8	21
	In vitro	13	14	8	25	9	25
<i>C. asiatica</i>	In vivo	12	11	11	18	10	15
	In vitro	10	9	13	23	20	21
<i>E. hirta</i>	In vivo	7	12	10	24	6	15
	In vitro	8	18	12	28	8	19

The table indicates that the antimicrobial effectiveness of callus cultures of *E. hirta* with inhibition zone of 18 mm was found highest against *C. gramineum* and in *O. tenniflorum* with inhibition zone of 28 mm was found maximum against *K. pneumoniae*. On the contrary, the *in vivo* analysis of antimicrobial efficacy of *O. tenniflorum* with a 16 mm inhibition zone was found to be higher against *P. chrysogenum* and *E. hirta* with a 24 mm inhibition zone was found to be higher against *K. pneumoniae*. The ethanolic extract generally exhibits a high degree of antimicrobial effectiveness, which appears to support the traditional therapeutic claims of these herbs (Dwivedi *et al.*, 1998). The most promising plants are those for which the extract should marked or significant activity. These results suggest the presence of either a good antibacterial potency or a high concentration of active ingredient in the extract. This

antimicrobial activity would support the traditional therapy for infections whose symptoms might involve fungi and bacteria. The results obtained indicated the existence of antimicrobial compounds in the crude ethanolic extracts of *O. tenniflorum*, *C. asiatica*, *E. hirta* and this shows that the ethanolic extract of these exhibits a broad spectrum of microbial inhibition.

4. Conclusions

This study is a preliminary assessment of antimicrobial activity of the plants. It indicates that several plants have the potential to generate novel metabolites. Crude extracts demonstrating antimicrobial activity could lead to the discovery of novel antimicrobial agents. Plants demonstrating a broad spectrum of activity could help discover new chemical classes of antibiotics that could serve as selective agents for the maintaining

animal or human health and provide biochemical tools for the study of infectious diseases.

Acknowledgements

We are grateful to the Institute of Environmental Science (IES), University of Rajshahi, Department of Pharmacy, University of Rajshahi, Department of Microbiology and Department of Biochemistry and Molecular Biology, GonoBishwabidyalay and BCSIR laboratories, Rajshahi, Bangladesh where all of the works were carried out.

References

- Dygerak M, Bagecy, E and Alma MH. 2002. *Pharmaceutical Biol.*,**40(6)**:425-428.
- Ibrahim SA, Dharmavaram SR, Seo CW and Shahbazi G. 2002. *Intern. J. Food Safety.*,**2**:6-8.
- Bassam A, Ghaleb A, Naser J, Awni A and Kamel A. 2006. *Turk J. Biol.*, **30**:195-198.
- Sasidharan VK, Krishnakumar T and Manjula, C.B. 1998. *Philippine J. sci.*,**127**: 65-71.
- Semra L, Filiz S, Ferda C, Cansu F and Zerrin FE. 2006. *Turk J. Biol.*,**30**:149-152.
- Suffredini IB, Paciencia MLB, Varella AD and Younes RN. 2006. *The Braz. J. Infectious Diseases.*,**10(6)**: 400-402.
- Kubmarawa D, Ajoku GA, Enwerem NM and Okorie DA. 2007. *Afr. J. Biotechnol.*, **6(14)**:1690-1696.
- Gould JC and Bowie JH. 1952. *Edinb. Med. J.*,**5**:59, 178.
- Ahmad I, Mahmood Z and Mohammad F. 1998. *J. Ethnopharmacol.*,**62**:183-193.
- Faria MHG, Carvalho TG, Rabenhorst SHB, Sidrim JC and Moraes MO. 2006. *Braz. J. Biol.*,**66(4)**: 1133-1135.
- Dwivedi SK and Singh KP. 1998. *Flavour and Fragrance J.*,**13**: 397-399.
- Mohammed AH. 2019. Importance of medicinal plants. *Res. Pharmacy Health Sci.*, **5(2)**:124-125.
- Kumar PM, Surya CM and Gopi VP. 2017. Identification of ayurvedic medicinal plants by images processing of leaf samples. Proceedings of the 3rd International Conference on Research in Computational Intelligence and Communication Networks (ICRCICN), IEEE., 231-238 Google Scholar.
- Wu SG, Bao FS, Xu EY, Wang YX, Chang YF and Xiang QL. 2007. A leaf recognition algorithm for plant classification using a probabilistic neural network. Proceedings of the IEEE International Symposium on Signal Processing and Information Technology, IEEE., 11-16. [View in Scopus](#) Google Scholar.
- Medical Botany. 2023. United States department of agriculture. <https://www.fs.usda.gov/wildflowers/ethnobotany/medicinal/index.shtml>.

